

UNITED STATES ANTARCTIC PROGRAM

# Field First Aid







This Manual was prepared for the U.S. National Science Foundation Office of Polar Programs (NSF/OPP) by Antarctic Support Contract (ASC) field support personnel and grantees. It brings together decades of first-hand field experience in Antarctica with the United States Antarctic Program (USAP).

Suggestions and corrections are encouraged and should be sent to [DEN-FieldSafety@usap.gov](mailto:DEN-FieldSafety@usap.gov).

Written and compiled by Cara Ferrier, Meghan Walker, and Kaija Webster.  
Edited and updated by Kate Koons, Al Davies, Jessie Crain and Sara Eckert.  
Design and layout by Hannah Halka.

Cover photo by Ben Urmston.

# Table of Contents

## First Aid

<b>Introduction</b> .....	8
<b>Hygiene</b> .....	8
<b>Sprains and Strains</b> .....	8
<b>Sprain and Strain Signs and Symptoms</b> .....	8
<b>Sprain and Strain Treatment</b> .....	9
<b>Immobilize the Joint</b> .....	9
<b>Bleeding and Wounds</b> .....	10
External Bleeding and Wound Treatment .....	10
Impaled Objects .....	10
<b>Carbon Monoxide Poisoning</b> .....	10
Carbon Monoxide Poisoning Signs and Symptoms .....	11
Carbon Monoxide Poisoning Treatment .....	11
<b>Hypothermia</b> .....	12
Hypothermia Prevention .....	12
Hypothermia Signs and Symptoms .....	12
Hypothermia Treatment .....	13
<b>Frostbite</b> .....	13
Frostbite Prevention .....	13
Mild Frostbite Signs and Symptoms (Pre-Thaw) .....	14
Full-Thickness Frostbite Signs and Symptoms (Pre-Thaw) .....	14
Full-Thickness Frostbite Signs and Symptoms (Post-Thaw) .....	14
Frostbite Treatment .....	15
Superficial Frostbite Treatment (Frost Nip) .....	15
Partial-Thickness Frostbite Treatment .....	15
Full-Thickness Frostbite Treatment .....	15
<b>Immersion Foot</b> .....	16
Immersion Foot Prevention .....	16
Immersion Foot Signs and Symptoms .....	16
Immersion Foot Treatment .....	17
<b>Altitude Sickness</b> .....	17
Altitude Sickness Prevention .....	17
Altitude Sickness Signs and Symptoms .....	18
Altitude Sickness Treatment .....	19
HACE Treatment .....	20
HAPE Treatment-URGENT .....	20
Acute Mountain Sickness (AMS) - Mild to Moderate .....	21
High Altitude Cerebral Edema (HACE) .....	21
High Altitude Pulmonary Edema (HAPE) .....	21

Treatment for mild to moderate AMS .....	22
Treatment for HACE/HAPE .....	22
<b>Eye Injuries</b> .....	22
Tent Eye .....	22
Snow Blindness .....	22
Snow Blindness Signs and Symptoms .....	22
Snow Blindness Treatment .....	22
Snow Blindness Prevention .....	23
<b>Skin Injuries</b> .....	23
Sunburn and Windburn .....	23
Sunburn and Windburn Prevention .....	23
<b>Dental Health</b> .....	23
Oral Hygiene .....	23
<b>Controlled Medications</b> .....	24
Issue of Restricted Drugs .....	24
<b>Chain of Custody</b> .....	24



Continental Field Manual

# First Aid

## Introduction

This First Aid Manual describes some of the medical issues that may be encountered in an Antarctic field environment. This is not an exhaustive manual, but it will serve as a guide for those with limited medical background to help treat their companions. It should be read in advance to help field team members recognize dangerous situations and help prevent injuries from occurring or becoming more serious if they do occur.

Antarctica is an inherently dynamic environment. Participants are often dehydrated, mildly hypothermic, and sleep deprived. This can lead to an increase in accidents. It is essential that all team members slow down, assess each task, and ensure the safety of the team.

There are a variety of medical kits provided to teams depending on their activities, locations and needs. Team members should familiarize themselves with the contents of the kit before there is a need to use it.

## Hygiene

Occasionally, people deploying to the deep field use the remote environment as an excuse to abstain from normal hygiene and sanitation. Extreme cold temperatures, lack of running water, and communal living make bathing, brushing teeth, and basic hygiene a chore. However, it is important to continue with a normal hygiene routine to avoid painful and distracting issues that can occur in the absence of cleanliness, such as dental abscesses, gum pain, yeast infections, skin rashes, cracked skin and trench foot.

Wet wipe and sponge baths are the norm. If wet wipes are preferred, personnel should bring enough for the anticipated time in the field. Blanket partitions can be set up in large tents to create a semi-private space.

## Sprains and Strains

Sprains and strains are the most common injuries in Antarctica. People must work carefully, thoughtfully and deliberately to avoid them. A sprain is an injury that involves tearing the ligaments that help keep joints intact. A strain involves overstretching a muscle.

## Sprain and Strain Signs and Symptoms

Sprains and strains will manifest as pain at the site of the injury that may radiate outward. There will also be swelling and discoloration.



## Sprain and Strain Treatment

- R Rest:** Stop activity, make the patient comfortable, and set up shelter if necessary.
- I Ice:** Cool down the affected area with water, snow or ice for approximately 15 minutes. Don't apply directly to the skin.
- C Compression:** Wrap the affected area with an elastic bandage.
- E Elevation:** Keep the affected limb raised to reduce swelling.

## Immobilize the Joint

### *Fingers*

Place gauze between the injured finger and the uninjured finger next to it. Buddy tape the fingers in two places but not over a joint, so the fingers can be flexed and extended.

### *Wrist*

If swollen, splint it using a SAM™ splint or materials on hand. If it is not swollen, or when the swelling subsides, bandage it from the hand to below the elbow with an elastic (ACE™) bandage. The patient should exercise the fingers, elbow and shoulder regularly.

### *Knee*

If it is very swollen, suspect a more serious injury. With the knee in a neutral position (slightly bent, between full extension and flexion) wrap a thick layer of cotton wool around the leg from mid-calf to mid-thigh. Apply a SAM™ splint on the back of the leg to keep it in position and hold the splint in place with an elastic bandage.

### *Ankle*

Remove the boot. Place the foot in a position of comfort. Using an elastic or tape bandage, wrap from the toes to just below the knee, keeping the foot up and covering all the skin. If the ankle is very swollen, especially on both sides, then a more severe injury should be suspected.

*Caution: Do not wrap bandages too tightly. Toes must remain warm and pink and have feeling.*

# Bleeding and Wounds

## External Bleeding and Wound Treatment

Stop the bleeding with a gloved hand or finger(s) using well-aimed, direct pressure. Once bleeding has been controlled, clean the wound with mild soap and potable water, or use Hibiclens® or povidone iodine. Remove any obvious debris with tweezers or forceps and irrigate the wound copiously with high pressure irrigation. If high pressure irrigation is not available, continue to flush the wound until no debris is seen. Never use alcohol to clean a wound. Once the wound is clean, apply an antibiotic ointment, cover the wound with a clean dressing (sterile, if possible) that extends well beyond the edges of the wound, and tape it in place.

Small wounds can be taped together with Steri-strips® or Band-Aids®. Dry the edges of the wound, squeeze together, and stick the Band-Aid® across the wound. Several may be needed to hold the cut together.

All wounds should be cleaned twice a day and a new, clean dressing applied. If a wound is contaminated and there is concern about infection, medical personnel should be notified.

## Impaled Objects

Impalements to the head, face, trunk and other core parts of the body are usually left in place. Primary focus should be on bleeding control and secondary efforts to stabilize the object in place with sterile bulky dressing. Minimizing movement of the impalement will help to reduce subsequent damage. In some cases when the impalement is from a larger object, consider reducing the size of the object to facilitate evacuation. Impalements to the eyes get special attention. No objects impaled in an eye should be removed. Instead, both eyes should be patched and padding should be placed around the injured eye to support it and decrease movement until it can be safely removed in a controlled environment.

Wilderness medicine guidelines suggest that impalements to the extremities can be removed to but the ideal scenario is to only remove objects that cannot be stabilized, will fall out easily or prevents transportation. If needed because of one of the above criteria, prior removal of the impaled object can help. Proper wound management will help minimize the risk of infection. In some cases, extremity impalements remain stabilized in place.

## Carbon Monoxide Poisoning

Carbon monoxide (CO) can be produced by burning anything containing carbon, including fuel in open flames, gas cookers, or engines. CO poisons by attaching itself to the hemoglobin in the blood. It does so about 200 times more readily than oxygen, easily displacing inhaled oxygen. When enough hemoglobin is compromised, the remainder cannot carry sufficient oxygen to the rest of the body. Oxygen starvation of the brain will cause permanent damage, even if the patient is revived. Furthermore, the toxicity of CO increases with altitude.

## **Carbon Monoxide Poisoning Signs and Symptoms**

Often there are none. However, the following may occur:

### ***Early Symptoms***

- Slight headache
- Dizziness
- Nausea
- Fatigue

### ***Progressed Symptoms***

- Shortness of Breath
- Confusion
- Panting
- Chest Pains
- Dimming of vision/blurry vision
- Loss of consciousness
- Ringing in the ears

### ***In Later Stages/More Rare Symptoms:***

- Patients with dark skin can appear as dusky, darker or even purplish.
- Patients with white skin will become pink to cherry red in the face, though the red and yellow polar tents will make it difficult to notice any skin color change.
- Unconsciousness and death are often rapid.

## **Carbon Monoxide Poisoning Treatment**

If carbon monoxide poisoning is suspected:

1. Immediately move the patient to fresh air or to an uncontaminated tent.
2. Provide the patient with 100% oxygen, if available.
3. Contact Medical and describe the incident and symptoms.
4. Keep the patient quiet and resting for at least eight hours. Early exertion may cause cardiac arrest.
5. If breathing stops, commence cardio-pulmonary resuscitation (CPR).

# Hypothermia

Hypothermia occurs when a person's core temperature is reduced to a level where normal brain and body functions are impaired. Hypothermia progressively affects a person's judgment, perception and coordination.

Wind greatly increases the chilling effect of cold. The faster the air moves, the more heat it can drag away. The cooling effects of air can be seen by referring to the wind chill chart in the reference section.

Wet clothes, from sweating, marine dampness, or precipitation, also cause chilling. Finally, fatigue reduces a person's ability to protect himself or herself, and it diminishes the physiological capacity to thermoregulate and maintain a proper core temperature.

## Hypothermia Prevention

Hypothermia is prevented by wearing the proper clothing and by supporting and regulating the body's heat production. Proper nutrition and hydration help prevent hypothermia, and adequate rest is critical. Exhaustion promotes the onset of hypothermia and precedes its development in almost all cases. The tendency to "press on" has led to many unnecessary deaths.

A layered clothing system should be employed, where layers can be added or removed as needed. Field team members should not allow themselves to get either cold or hot and sweaty.

## Hypothermia Signs and Symptoms

Hypothermia manifests in three stages:

### *Mild*

This stage includes shivering and personality changes. A person may become withdrawn, apathetic, or irritable. There is a loss of fine motor control. Field party members should always be on the alert for a team member displaying the "umbles": stumbles, mumbles, fumbles and grumbles.

### *Moderate*

At this stage, hypothermia progresses to violent shivering, altered mental states, and disorientation. Moderate hypothermia also manifests as a loss of gross motor skills, such as balance and coordination (ataxia).

### *Severe*

In this stage, shivering stops and the level of responsiveness drops. A person becomes unresponsive and may appear dead, with slow and weak pulse and

respiration rates. An individual will appear cold and blue, and he or she may have associated frostbite. Cardiac arrest is possible.

## Hypothermia Treatment

The essential and immediate treatment for hypothermia is to prevent further heat loss by insulating the body. If any member of a field party shows signs of developing hypothermia, the individual must be moved into shelter immediately.

Mild hypothermia may be turned around quickly. A person with this condition should be:

- Helped into additional clothing layers and fed quick-energy carbohydrates and warm, sweet drinks, such as hot chocolate or warm electrolyte beverages.
- Encouraged to run in place or perform another exercise.
- Provided dry clothes, if necessary, and external heat sources, such as hot pads or water bottles filled with warm fluid.

If moderate to severe hypothermia is suspected, contact Medical immediately. The patient should be placed in a hypothermia wrap, which is a bundle made of sleeping bags and reflective sheeting, with warm heat sources on the patient's neck, armpits and groin. Body-to-body rearming in a sleeping bag is of limited usefulness and may result in two cold people.

## Frostbite

Frostbite is freezing of body tissue. Areas most at risk are the extremities and exposed skin (ears, nose, face). Factors that lead to frostbite are:

- Previous frostbite injury
- Cold temperatures and wind
- High altitude
- Overexertion (fatigue and dehydration)
- Touching metal or super-cooled liquid fuel
- Poor circulation
- Constrictive clothing or footwear
- Underlying medical problems
- Hypothermia

## Frostbite Prevention

Frostbite is almost always avoidable. A buddy system should be established to observe any whitening on the face or ears of a companion. If any whitening or tingling of the face, ears, feet, or hands occurs, these areas should be warmed

immediately. Socks and boots should fit snugly, with no points of tightness. Liner gloves should be worn so that skin is never exposed when performing work that can't be done in heavy gloves.

If the body's core gets too cold, the body will restrict blood flow to the extremities to prevent damage to internal organs. This restriction increases the chance of frostbite. Strenuous exercise should be avoided in extreme cold, particularly at high altitudes. Very cold air brought too rapidly into the lungs will chill the body's core. Perspiration under conditions of extreme cold should be avoided. Perspiration evaporates, chilling the body.

Plenty of food should be consumed to produce maximum output of body heat. Food items in cold weather should tend toward quick energy first, such as fats and carbohydrates, and then proteins. In addition, personnel should drink two to three liters of water per day to stay hydrated.

Avoid the following, which can promote the occurrence of frostbite:

- Smoking
- Alcohol
- Excessive coffee and tea drinking
- Excessive fatigue
- Improper or inadequate eating habits
- Unnecessary medication
- Exposure to fuel, especially on bare skin

### **Mild Frostbite Signs and Symptoms (Pre-Thaw)**

There is an uncomfortable sensation of coldness, followed by numbness and skin anesthesia. In superficial frostbite (frost nip), white skin turns red, then pale or waxy-white. In dark skin individuals, the skin will appear lighter than their normal skin tone. In partial thickness frostbite, the skin becomes cold and frozen on the surface, but remains soft and pliable when gently pressed.

### **Full-Thickness Frostbite Signs and Symptoms (Pre-Thaw)**

The skin is waxy-white. Toes and fingers become solid like a piece of chicken taken from the freezer. They feel wooden, and the skin cannot be rolled over the bone.

### **Full-Thickness Frostbite Signs and Symptoms (Post-Thaw)**

The entire hand or foot swells, which limits the mobility of the injured toes or fingers. Blue, violet or grey (the worst) discoloration appears. After two days, the patient suffers severe throbbing and shooting pains. Huge blisters form, usually between the third and seventh day. These usually dry up, blacken, and

slough off, leaving an exceptionally sensitive thin, red layer of new skin.

## **Frostbite Treatment**

Frostbite should not be rubbed, as this will cause additional tissue damage from the ice crystals within. Treatment in the field for anything beyond superficial frostbite is full of challenges and additional risk for the patient. Prevention is paramount!

### **Superficial Frostbite Treatment (Frost Nip)**

Superficial frostbite can be treated effectively in the field. If noticed promptly, it can usually be treated by the firm, steady (no rubbing) pressure of a warm hand, or by blowing onto it with warm breath. Superficially frostbitten feet are best treated by removing the patient's footwear the moment there is any suspicion of danger and rewarming the feet immediately. After warming is complete, the affected feet should be covered with dry socks. If footwear is replaced, it should be done loosely to ensure adequate circulation so that warmth is maintained.

### **Partial-Thickness Frostbite Treatment**

Partial-thickness frostbite of a small body area should be reheated in water that is between 42°C and 43°C (107°F and 109°F). Water at higher temperatures can burn the skin. The injury should then be treated to prevent infection, bandaged for protection, and kept warm. Refreezing must be prevented, as this will cause major additional damage. Medical personnel should be consulted if necessary.

### **Full-Thickness Frostbite Treatment**

Because of limited resources in the field, full-thickness frostbite is a major medical emergency. Medical personnel should be contacted immediately for consultation and to discuss evacuation plans. Rewarming should not be attempted in the field if there is any possibility that the affected part may become refrozen. In such cases, the affected part must be kept frozen until it can be re-warmed rapidly under controlled conditions.

If rewarming is recommended, remove jewelry (especially rings) if possible. Immerse the injured part in 107°F to 109°F (42°C to 43°C) water, continually adding water and stirring it to maintain a constant temperature, until the digital tips (ends of fingers or toes) turn pink or burgundy red. This takes approximately 20 minutes to one hour. When adding water, take care that the water is not more than 111°F (44°C) and is not poured directly over the injured body part.

Significant pain, swelling, and blistering will develop after re-warming. Do not puncture the blisters, and do not allow the injury to refreeze.

Protect the thawed injury with sterile, soft, fluffy dressings. Separate toes and fingers with cotton wool. Wrap the whole part lightly with gauze bandages. Do not change dressings unless they get dirty, and never rub the skin. Keep the patient and the injured body part warm. Pain medication will be needed, and medical personnel will advise on specific type and dose. In addition:

- Elevate the injured limb(s).
- Commence antibiotic treatment, per instruction from the medical team.
- Keep the patient absolutely still and lying down.
- Evacuate to a medical facility as quickly as possible.

## Immersion Foot

Immersion foot, or trench foot, is a medical condition caused by prolonged exposure to cold, damp and unsanitary conditions.

### Immersion Foot Prevention

Feet should be kept warm and dry by wearing protective footwear, and they should be checked frequently during wet and cold conditions. Footwear should not be constrictive, and it should be cleaned and dried at every opportunity. In the field, extra pairs of dry socks should be carried next to the abdomen under the shirt. Wet socks can be dried by placing them next to the abdomen, either inside or outside the shirt.

If feet get wet, they should be dried as soon as possible. They can be warmed by the hands. Foot powder should be applied and dry socks put on. If it is necessary to wear wet socks and footwear for any length of time, then the feet should be exercised at regular intervals by wriggling the toes and bending the ankles.

### Immersion Foot Signs and Symptoms

The area becomes cold, swollen, waxy-white and mottled with burgundy-to-blue plotches. The skin becomes numb, deep sensation is lost, and movement of the affected area becomes difficult.

If allowed to continue untreated, the area becomes red, hot, and swollen, and blisters appear. The victim experiences constant throbbing and a burning sensation. Skin numbness is aggravated by heat and relieved by cold.



## Immersion Foot Treatment

Remove wet footwear. Gently and rapidly rewarm the affected foot by immersing it in warm water (about 104°F or 40°C). Once the foot is warmed, dry it completely and elevate it in a warm space. Swaddle it with clean bandages or cloth to keep it warm and clean.

The injury must not be rubbed or massaged. Blisters should be kept clean and dry. Do not apply ointments. Two 200 mg ibuprofen tablets every four hours may be administered for pain, if required. Evacuate the victim to a medical facility.

## Altitude Sickness

Altitude sickness also called acute mountain sickness, or AMS is caused by the body not adapting to the reduced availability of oxygen at altitude (as elevation increases, the partial pressure of oxygen decreases). The higher the altitude, the more common AMS becomes. Symptoms may range from minor lethargy to a coma, and death may result, so any symptom must be treated with caution. Anyone can be affected by altitude sickness, regardless of age, fitness level, or previous experience at altitude. Healthy individuals may experience symptoms as low as 2,500 meters (8,000 feet). Beyond 3,000 meters (9,840 feet), 75% of people will experience some level of AMS. The symptoms usually start 12 to 24 hours after arrival at altitude and begin to decrease in severity around the third day. Mild AMS does not interfere with normal activity, but anyone experiencing symptoms should communicate this to others so the person can be monitored.

Many work sites in Antarctica, such as NSF Amundsen-Scott South Pole Station and Mount Erebus, are at high altitude. The Fang Glacier acclimatization camp on Mount Erebus is at 2,900 meters (9,500 feet), with South Pole Station only slightly lower. Since the polar atmospheric effect raises the pressure altitude 10% to 15% above actual elevation, both of those locations will feel like 3,200 meters (10,500 feet) or more, increasing the risk of AMS.

AMS risk is also increased by rapid ascent to altitude (e.g., by air-craft), so team members must factor sufficient time for acclimatization into their schedule when flying to a high-altitude site. Minimum work should be planned for the first few days.

## Altitude Sickness Prevention

It is common to fly to altitude in Antarctica, thus increasing the risk of

altitude sickness. For those traveling to Mout Erebus, which is 3,794 meters (12,450 feet) above sea level, spending at least two nights acclimatizing on the Fang Glacier seems to reduce altitude-related problems. Unfortunately, it is usually not possible to have acclimatization camps for polar plateau work. For those traveling to the plateau, bring altitude medicine (if medical personnel advise doing so), and pack Gamow bags and oxygen, if possible. In addition, taking the following steps can reduce the incidence of AMS:

- Do not overexert upon arrival at altitude. Take it easy for the first three days.
- Avoid depressant drugs like sleeping pills and narcotics.
- Avoid alcohol and tobacco.
- Get plenty of sleep.
- Stay hydrated.
- Consider taking altitude medication, such as Diamox®. Discuss this with the medical team beforehand.

Above all, adjust expectations of how much work can be completed in the first few days at altitude. Team members should check in on each other constantly and let each other know how they are doing. Withholding information could be dangerous and is poor expedition behavior. If a team member is still not feeling better after 72 hours, follow up with the medical team.

## **Altitude Sickness Signs and Symptoms**

### ***Mild to Moderate AMS***

Most people arriving at altitude will see their breathing rate increase immediately. Other mild symptoms include headache, nausea, fatigue and lack of appetite. The average time to recover from mild AMS is approximately three days. Full acclimatization may take two months. However, a small number of individuals are unable to acclimatize at all. Neither prior experience at altitude nor a high level of fitness exclude a person from contracting altitude-related illnesses.

### ***Moderate AMS***

Moderate AMS will manifest as a more severe headache (which is not relieved by medication), increased nausea and vomiting, increased lethargy, loss of appetite, light-headedness, disturbed sleep, shortness of breath (even while resting), and decreased coordination. Normal activity becomes difficult. Though these symptoms may be due to other causes, it is wise to assume they are due to AMS until proven otherwise. The only treatments for moderate AMS are advanced medication or immediate descent to a lower altitude, with the latter being the preferable option. Without treatment, moderate AMS

could become more severe.

### **Severe AMS**

Severe AMS may manifest as High-Altitude Cerebral Edema (HACE) or High-Altitude Pulmonary Edema (HAPE), or both. Signs and symptoms of severe HACE are loss of muscular coordination (ataxia), decreased mental status (e.g., confusion, coma), severe headache, weakness, and vomiting. It appears that persons who have had HACE in the past are more susceptible to developing it again.

### **HACE**

HACE is a potentially fatal condition. Altered consciousness and loss of coordination are the base markers for HACE. The individual will be unable to take care of basic needs i.e., eating and dressing. Within a day of losing coordination, HACE victims slip into a coma. Without proper medical care, death will result.

### **HAPE**

HAPE is the most common altitude-related cause of death. Early signs are decreased exercise performance and increased recovery time. Specific HAPE signs and symptoms are dry cough, shortness of breath at rest, a gurgling/crackling noise heard in chest during breathing, and pale or blue color to skin and nail beds (cyanosis). In the late stages, a wet, productive cough will be present. As with HACE, people who have had HAPE in the past have a high likelihood of developing it again. Both the Lower Hut on Mount Erebus and South Pole Station have seen more than a few cases of HAPE over the decades.

## **Altitude Sickness Treatment**

Acetazolamide (Diamox®) may be used as a preventive measure before going to altitude. It is a diuretic and respiratory stimulant that accelerates the body's acclimatization. With the consent of medical personnel, team members should begin taking it several days before ascent.

Dexamethasone (Decadron®) is no longer used as a preventive measure before ascending to altitude because prophylactic use has been found to diminish its efficacy. Decadron may be used prophylactically in very limited circumstances, but each case requires a full medical evaluation and a risk/benefit discussion with the patient, who must understand that such use will likely make it less effective in reducing altitude-induced, life-threatening swelling in the brain.

Mild to moderate AMS requires rest, medication for headaches and nausea, hydration, proper nutrition, and supplemental oxygen, if available. If AMS occurs, the best treatment is rapid descent. Contact medical personnel

to discuss the severity of the case and recommendations for evacuation. Supplemental oxygen is helpful, in addition to using Diamox® and/or Decadron®.

A Gamow bag is a portable hyperbaric chamber that simulates rapid descent. It has saved hundreds of lives at altitude worldwide since its introduction in 1990. If descent isn't possible because of flight delays or because team members cannot take the victim down themselves, a Gamow bag is the best asset to have. The elevation at which teams will be working will determine if the members need to be trained in Gamow bag operation and issued one for the field.

## **HACE Treatment**

Early, simple tests (similar to roadside sobriety tests) can be conducted to look for the loss of coordination that is the hallmark of HACE. Individuals may be asked to stand with eyes closed and arms extended to the side, then asked to touch their nose. Also, with eyes closed, they may be asked to walk forward, heel to toe. Be sure that someone is ready to catch them if they stumble. However, differentiating between moderate AMS and the initial stages of HACE may be difficult. Assume the worst and treat for HACE.

Severe AMS/HACE requires an urgent call to the medical team. Evacuation is likely. If immediate descent is not possible, the patient should be placed in a Gamow bag. Provide the patient with oxygen. Medical personnel may also prescribe Diamox® and Decadron®.

## **HAPE Treatment-URGENT**

If HAPE is suspected, contact the medical team immediately to request an evacuation. Immediately descend 600 to 1,200 meters (2,000 to 4,000 feet), if possible. Use a Gamow bag while waiting for evacuation and provide the patient with oxygen. Nifedipine may be helpful. The medical team can authorize administration of this medication.

### ***To Prepare***

- Drink at least three liters of water a day, starting three days before flying to altitude.
- Attend altitude orientation to learn the risks of high altitude and benefits of altitude acclimatization medication.
- Get good rest and eat a high carb diet. Sleep for 7-9 hours.
- Avoid alcohol for 48 hours before ascending to altitude. Alcohol increases dehydration and decreases respiration rate.
- Quit smoking. Smoking impacts acclimatization and lung function, and

carbon monoxide can build up in blood.

- Check in with the Medical lead so he or she can help monitor the acclimatization process.

### ***To Avoid***

- Avoid drinking alcohol before leaving and during the first 72 hours at altitude.
- No physical overexertion, at work or while exercising.
- Do not smoke, so your body can adjust easier to long-term, altitude acclimatization.
- DO NOT ignore your headache and other symptoms of Acute Mountain Sickness (AMS) during the first 24-72 hours.
- DO NOT put yourself and others in USAP at risk by not taking appropriate precautions and not understanding the severity of AMS and other altitude-related sicknesses.

Do not let AMS develop into the life-threatening conditions of High Altitude Pulmonary Edema or High Altitude Cerebral Edema.

“Altitude Sickness” is a common response that the body has from rapid ascent to altitude, that can become more serious in some individuals. Flying from McMurdo Station (sea level) to the South Pole (9,306 feet) or other high-altitude locations put you at risk. Acute Mountain Sickness that is ignored can progress and become life threatening. The following conditions can usually be prevented by early detection and monitoring for worsening signs and symptoms.

## **Acute Mountain Sickness (AMS) - Mild to Moderate**

### ***Signs and Symptoms***

HEADACHE with any or all of the following:

- Shortness of breath with mild exertion.
- Lack of energy and /or appetite (feel bad).
- Inability to sleep or restless sleep.

## **High Altitude Cerebral Edema (HACE)**

### ***Signs and Symptoms***

- AMS with severe headache (severe AMS).
- Altered mental status: disoriented, dizzy, and/or personality changes.
- Unconsciousness.

**THIS IS A MEDICAL EMERGENCY!**

## High Altitude Pulmonary Edema (HAPE)

### *Signs and Symptoms*

(May not have other signs and symptoms of AMS).

- Shortness of breath at rest.
- Inability to work or function normally.
- Cough that worsens and eventually becomes productive (wet).
- Pale coloring, increased heart rate and labored breathing.

**THIS IS A MEDICAL EMERGENCY!**

### **Treatment for mild to moderate AMS**

- Medication for headache.
- Rest, hydration and nutrition.
- Supplemental oxygen.

### **Treatment for HACE/HAPE**

- Evacuation (aircraft) to lower altitude ASAP.
- Oxygen.
- Gamow Bag.
- Medication from Medical or field team medical lead.

## Eye Injuries

### **Tent Eye**

Antarctica's extreme low humidity may cause the film of tears protecting the eye to dry up, making the cornea susceptible to damage from stove fumes in the tent. The condition can be treated by applying Chlorsig® ointment to the eye when it occurs and before going to sleep.

### **Snow Blindness**

Snow blindness is caused by ultraviolet (UV) light burning the eyes. The danger of snow blindness is greatest not on clear, bright days but on dull, cloudy (whiteout) days, when crystalline snow mist is present. There is no warning that damage has been done until the symptoms begin to appear two to twelve hours after exposure.

### **Snow Blindness Signs and Symptoms**

Snow blindness manifests as intensely painful, red, watering eyes that are sensitive to light. The victim will also feel as though there is grit in the eyes.

## **Snow Blindness Treatment**

A single episode of snow blindness may last up to five days, even while being treated. The eyes should be rested for at least 24 hours. That means closing them and covering them with a non-fluffy pad. If the temperature is above freezing, a cold compress may be placed over the affected eyes to relieve pain. Medical should be contacted for treatment recommendations and possible medications. Medical personnel may recommend providing the victim two 200mg tablets of ibuprofen every four hours, as required, or putting Chlorsig® ointment on the eyes every three hours.

## **Snow Blindness Prevention**

This condition must be avoided, as it is a crippling injury that may seriously delay a field party. Team members should wear dark, UV-protective glasses or goggles with the appropriate lenses (not yellow) at all times when in the field, especially on overcast days.

## **Skin Injuries**

### **Sunburn and Windburn**

Direct exposure to the sun, especially when it is very windy or the body is wet with sweat, can result in a sunburn and chaffed skin. Because the Antarctic air is cleaner and thinner, there is greater ultra-violet penetration, so sunburn can occur even on overcast days. If sunburn occurs, apply aloe vera gel to the burn and provide the victim 400 mg of ibuprofen every four hours, as necessary, to relieve pain.

### **Sunburn and Windburn Prevention**

Prevent sunburn by applying sunscreen ChapStick® to the lips and regular sunscreen to other areas of exposed skin. Covering the face with a balaclava will prevent both sunburn and windburn to this frequently exposed area.

## **Dental Health**

### **Oral Hygiene**

Oral hygiene can be inconvenient in the field, but it is just as important as bodily hygiene. Failure to maintain good oral hygiene may result in tooth decay (especially around the edges of fillings) and gingivitis. Ideally, teeth must be brushed after every meal, with snow if no water is available. Use toothpicks or waxed dental floss to clean gaps between the teeth that are hard to clean with the brush.

## Controlled Medications

### Issue of Restricted Drugs

The McMurdo Station clinic issues a field medication kit containing over-the-counter, prescription, and controlled (restricted) medications to each designated field party medical lead. This person is responsible for the field medication kit. The medical lead or any USAP participant must contact a station doctor for consultation and authorization before administering any prescribed medication. Always check for any known allergies before administering drugs.

### Chain of Custody

McMurdo Station clinic personnel will fill out a controlled drug Chain of Custody form and provide it to the field medical lead. The lead must account for all controlled substances when the kit is checked out, weekly, and when the kit is returned. If the lead departs before the end of the season, he or she must complete a new Chain of Custody form and count the medications before transferring the kit to another person. At the end of the season, the medical kit must be returned to the clinic.



